

# FDA Foodborne Illness-Causing Organisms in the U.S. WHAT YOU NEED TO KNOW

While the American food supply is among the safest in the world, the Federal government estimates that there are about 48 million cases of foodborne illness annually—the equivalent of sickening 1 in 6 Americans each year. And each year these illnesses result in an estimated 128,000 hospitalizations and 3,000 deaths.

The chart below includes foodborne disease-causing organisms that frequently cause illness in the United States. As the chart shows, the threats are numerous and varied, with symptoms ranging from relatively mild discomfort to very serious, life-threatening illness. While the very young, the elderly, and persons with weakened immune systems are at greatest risk of serious consequences from most foodborne illnesses, some of the organisms shown below pose grave threats to *all* persons.

ORGANISM	COMMON NAME OF ILLNESS	ONSET TIME AFTER INGESTING	SIGNS & SYMPTOMS	DURATION	FOOD SOURCES
<b><i>Bacillus cereus</i></b>	<i>B. cereus</i> food poisoning	10-16 hrs	Abdominal cramps, watery diarrhea, nausea	24-48 hours	Meats, stews, gravies, vanilla sauce
<b><i>Campylobacter jejuni</i></b>	Campylobacteriosis	2-5 days	Diarrhea, cramps, fever, and vomiting; diarrhea may be bloody	2-10 days	Raw and undercooked poultry, unpasteurized milk, contaminated water
<b><i>Clostridium botulinum</i></b>	Botulism	12-72 hours	Vomiting, diarrhea, blurred vision, double vision, difficulty in swallowing, muscle weakness. Can result in respiratory failure and death	Variable	Improperly canned foods, especially home-canned vegetables, fermented fish, baked potatoes in aluminum foil
<b><i>Clostridium perfringens</i></b>	Perfringens food poisoning	8–16 hours	Intense abdominal cramps, watery diarrhea	Usually 24 hours	Meats, poultry, gravy, dried or precooked foods, time and/or temperature-abused foods
<b><i>Cryptosporidium</i></b>	Intestinal cryptosporidiosis	2-10 days	Diarrhea (usually watery), stomach cramps, upset stomach, slight fever	May be remitting and relapsing over weeks to months	Uncooked food or food contaminated by an ill food handler after cooking, contaminated drinking water
<b><i>Cyclospora cayentanensis</i></b>	Cyclosporiasis	1-14 days, usually at least 1 week	Diarrhea (usually watery), loss of appetite, substantial loss of weight, stomach cramps, nausea, vomiting, fatigue	May be remitting and relapsing over weeks to months	Various types of fresh produce (imported berries, lettuce, basil)
<b><i>E. coli (Escherichia coli) producing toxin</i></b>	<i>E. coli</i> infection (common cause of “travelers’ diarrhea”)	1-3 days	Watery diarrhea, abdominal cramps, some vomiting	3-7 or more days	Water or food contaminated with human feces
<b><i>E. coli</i> O157:H7</b>	Hemorrhagic colitis or <i>E. coli</i> O157:H7 infection	1-8 days	Severe (often bloody) diarrhea, abdominal pain and vomiting. Usually, little or no fever is present. More common in children 4 years or younger. Can lead to kidney failure	5-10 days	Undercooked beef (especially hamburger), unpasteurized milk and juice, raw fruits and vegetables (e.g. sprouts), and contaminated water
<b>Hepatitis A</b>	Hepatitis	28 days average (15-50 days)	Diarrhea, dark urine, jaundice, and flu-like symptoms, i.e., fever, headache, nausea, and abdominal pain	Variable, 2 weeks-3 months	Raw produce, contaminated drinking water, uncooked foods and cooked foods that are not reheated after contact with an infected food handler; shellfish from contaminated waters
<b><i>Listeria monocytogenes</i></b>	Listeriosis	9-48 hrs for gastro-intestinal symptoms, 2-6 weeks for invasive disease	Fever, muscle aches, and nausea or diarrhea. Pregnant women may have mild flu-like illness, and infection can lead to premature delivery or stillbirth. The elderly or immunocompromised patients may develop bacteremia or meningitis	Variable	Unpasteurized milk, soft cheeses made with unpasteurized milk, ready-to-eat deli meats
<b>Noroviruses</b>	Variously called viral gastroenteritis, winter diarrhea, acute non-bacterial gastroenteritis, food poisoning, and food infection	12-48 hrs	Nausea, vomiting, abdominal cramping, diarrhea, fever, headache. Diarrhea is more prevalent in adults, vomiting more common in children	12-60 hrs	Raw produce, contaminated drinking water, uncooked foods and cooked foods that are not reheated after contact with an infected food handler; shellfish from contaminated waters
<b><i>Salmonella</i></b>	Salmonellosis	6-48 hours	Diarrhea, fever, abdominal cramps, vomiting	4-7 days	Eggs, poultry, meat, unpasteurized milk or juice, cheese, contaminated raw fruits and vegetables
<b><i>Shigella</i></b>	Shigellosis or Bacillary dysentery	4-7 days	Abdominal cramps, fever, and diarrhea. Stools may contain blood and mucus	24-48 hrs	Raw produce, contaminated drinking water, uncooked foods and cooked foods that are not reheated after contact with an infected food handler
<b><i>Staphylococcus aureus</i></b>	Staphylococcal food poisoning	1-6 hours	Sudden onset of severe nausea and vomiting. Abdominal cramps. Diarrhea and fever may be present	24-48 hours	Unrefrigerated or improperly refrigerated meats, potato and egg salads, cream pastries
<b><i>Vibrio parahaemolyticus</i></b>	<i>V. parahaemolyticus</i> infection	4-96 hours	Watery (occasionally bloody) diarrhea, abdominal cramps, nausea, vomiting, fever	2-5 days	Undercooked or raw seafood, such as shellfish
<b><i>Vibrio vulnificus</i></b>	<i>V. vulnificus</i> infection	1-7 days	Vomiting, diarrhea, abdominal pain, bloodborne infection. Fever, bleeding within the skin, ulcers requiring surgical removal. Can be fatal to persons with liver disease or weakened immune systems	2-8 days	Undercooked or raw seafood, such as shellfish (especially oysters)

For more information, contact: The U.S. Food and Drug Administration Center for Food Safety and Applied Nutrition  
 Food Information Line at 1-888-SAFEFOOD (toll free), 10 AM to 4 PM ET, Monday through Friday.  
 Or visit the FDA Web site at [www.fda.gov](http://www.fda.gov).